



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

years, and it is deemed to be no longer necessary to issue a daily report of quarantinable diseases.

No vessels cleared for United States ports.

FOREIGN AND INSULAR STATISTICAL REPORTS OF COUNTRIES AND CITIES—
UNTABULATED.

BRAZIL.—State of São Paulo, including the cities of *São Paulo*, *Santos*, and *Campinas*.—Week ended December 30, 1906. Estimated population, 286,000. Total number of deaths, 187, including whooping cough 2, enteric fever 2, measles 4, leprosy 2, and 17 from tuberculosis.

Pernambuco.—Two weeks ended December 31, 1906. Estimated population, 210,000. Total number of deaths, 278, including leprosy 1, plague 1, whooping cough 2, smallpox 48, and 39 from tuberculosis.

Rio de Janeiro.—Month of October, 1906. Estimated population, 912,900. Total number of deaths, 1,162, including diphtheria 2, enteric fever 6, bubonic plague 18, measles 1, whooping cough 5, beriberi 1, leprosy 1, yellow fever 3, and 268 from tuberculosis.

CUBA—*Cardenas*.—Months of November and December, 1906. Estimated population, 27,935. Total number of deaths, 114, including diphtheria 1, leprosy 1, and 18 from tuberculosis.

FRANCE—*St. Etienne*.—Two weeks ended January 15, 1907. Estimated population, 146,836. Total number of deaths, 198, including whooping cough 1, enteric fever 2, and 29 from tuberculosis.

GIBRALTAR.—Two weeks ended January 27, 1907. Estimated population, 27,385. Total number of deaths, 14. No deaths from contagious diseases reported.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 76 great towns in England and Wales during the week ended January 26, 1907, correspond to an annual rate of 18.2 per 1,000 of population, which is estimated at 16,024,458.

London.—One thousand six hundred and forty-seven deaths were registered during the week, including measles 17, scarlet fever 11, diphtheria 16, enteric fever 5, whooping cough 35, and 16 from diarrhea. The deaths from all causes correspond to an annual rate of 18 per 1,000. In Greater London 2,378 deaths were registered. In the "outer ring" the deaths included 4 from diphtheria, 1 from measles, 6 from whooping cough, and 2 from scarlet fever.

Ireland.—The average annual death rate represented by the deaths registered during the week ended January 26, 1907, in the 21 principal town districts of Ireland was 24.6 per 1,000 of the population, which is estimated at 1,117,547. The lowest rate was recorded in Armagh, viz, 6.9, and the highest in Kilkenny, viz, 44.2 per 1,000. In Dublin and suburbs 225 deaths were registered, including scarlet fever 1, enteric fever 1, whooping cough 6, and 41 from tuberculosis.

Scotland.—The deaths registered in 8 principal towns during the week ended January 26, 1907, correspond to an annual rate of 19.1 per 1,000 of the population, which is estimated at 1,812,171. The lowest rate of mortality was recorded in Leith, viz, 15.6, and the highest in Perth, viz, 26.9 per 1,000. The aggregate number of deaths registered from all causes was 662, including diphtheria 4, enteric fever 2, scarlet fever 7, typhus fever 1, measles 3, and 16 from whooping cough.

HAWAII.—Year ended June 30, 1906: *Hilo.*—Total number of deaths, 347, including bubonic plague 1, diphtheria 4, enteric fever 15, whooping cough 2, and 34 from tuberculosis.

Honolulu.—Population, 39,306. Total number of deaths, 880, including bubonic plague 22, diphtheria 6, enteric fever 25, and 134 from tuberculosis.

Island of Oahu (exclusive of *Honolulu*)—Total number of deaths, 281, including bubonic plague 2, diphtheria 1, enteric fever 12, whooping cough 3, and 34 from tuberculosis.

Island of Hawaii (exclusive of *Hilo*)—Total number of deaths, 518, including bubonic plague 3, diphtheria 9, enteric fever 35, measles 1, whooping cough 8, and 54 from tuberculosis.

Island of Kauai: Total number of deaths, 244, including diphtheria 3, enteric fever 8, and 25 from tuberculosis.

Islands of Maui, Molokai, and Lanai: Total number of deaths, 528, including diphtheria 1, enteric fever 13, and 57 from tuberculosis.

JAMAICA—Kingston.—Three weeks ended January 26, 1907. Estimated population, 52,065. Total number of deaths not reported. Two deaths from enteric fever reported.

MALTA.—Three weeks ended January 26, 1907. Estimated population, 205,059. Total number of deaths, 350, including diphtheria 6, enteric fever 1, and 1 from whooping cough.

ST. HELENA.—Five weeks ended January 12, 1907. Estimated population, 3,500. Total number of deaths, 3. No deaths from contagious diseases.

SWITZERLAND.—Week ended January 12, 1907. Reports from 18 cities, having an aggregate population of 867,056, show as follows: Total number of deaths, 326, including enteric fever 2, measles 21, whooping cough 1, and 37 from tuberculosis.

URUGUAY—Montevideo.—Month of November, 1906. Estimated population, 306,890. Total number of deaths, 430, including diphtheria 1, enteric fever 1, measles 16, scarlet fever 2, whooping cough 3, and 82 from tuberculosis.

WEST INDIES—Curacao.—Two weeks ended February 1, 1907. Estimated population, 31,600. Total number of deaths, 18. No contagious diseases reported.